

ON THE FREE BURNSIDE GROUP AS A FACTOR OF VON DYCK GROUP: A GEOMETRIC PERSPECTIVE

GIOVANNI MORENO* AND MONIKA EWA STYPA**

ABSTRACT. Von Dyck groups have a natural geometric interpretation which explains certain properties of free Burnside groups with two generators. We propose an algorithmic method to enumerate the Fuchsian von Dyck groups $D(n, n, n)$ which can be used to determine the finiteness of some Burnside groups.

CONTENTS

Introduction	2
Notation and conventions	4
1. Colored graphs and their morphisms	4
2. The colored graph of a group with a Borel-free set of generators	8
3. $2n$ -gonal tiling of constant-curvature surfaces and their 2-colors graphs	10
4. Triangle and von Dyck groups	11
5. Free Burnside groups	17
5.1. Finiteness of $B(2, 3)$: an algebraic proof	19
5.2. Finiteness of $B(2, 4)$: an algebraic proof	21
6. The kernel subgroup K_n	22
6.1. Finiteness of $B(2, 3)$: a geometric proof	23
7. Representing the graph of $B(2, n)$ on a quotient of a constant-curvature surfaces	24
7.1. Higher-order von Dyck groups?	27
References	28

2010 *Mathematics Subject Classification*. Primary: 05E18; 20F65; Secondary: 05C25; 05C15; 52C20; 51E99; 20F05; 22E40;

Key words and phrases. Group actions on combinatorial structures, Geometric group theory, Incidence Geometry, Graphs and abstract algebra, Tilings in 2 dimensions.

The authors are grateful to P. Longobardi and C. Sica for sharing their expertise in group theory. The first author is thankful to the Grant Agency of the Czech Republic (GA ČR) for financial support under the project P201/12/G028. The second author is grateful to the Ph.D. program of Salerno University for financial support.

INTRODUCTION

We study the relationship between two historic groups. One, denoted by $D(a, b, c)$, was introduced by Walther von Dyck at the turn of eighteen century (see, e.g., [25]). Another, denoted by $B(m, n)$, appeared few years later (1902) as William Burnside formulated his celebrated problem [5]. From a mere group-theoretic standpoint, there is a rather evident¹ though unexplored projection of $D(n, n, n)$ over $B(2, n)$. The geometric implications of this projection is the main concern of the present paper.

Depending on the value of n , the group $D(n, n, n)$ can be generated by orientation-preserving transformations of a tiling $\mathcal{T}_n$ of a constant curvature surface (denoted, from now on, by the symbol $\mathbb{S}$) by regular triangles. As opposed to von Dyck groups, the definition of a free Burnside group is rather abstract: $B(m, n)$ is the free object over a rank- m set, in the variety of groups determined by the equation $x^n = 1$. However, it is precisely the epimorphism between $D(n, n, n)$ and $B(2, n)$ which makes it possible to associate to the latter some of the geometric features of the former.

Intuitively, if $D(n, n, n)$ acts on the tiling $\mathcal{T}_n$ of $\mathbb{S}$, then all its factors, including $B(2, n)$, must act on some sort of “quotient tiling” of $\mathcal{T}_n$, with respect to the kernel, henceforth denoted by K_n , of the projection $D(n, n, n) \rightarrow B(2, n)$. Though formally correct, this perspective requires working with an orbifold of $\mathbb{S}$, thus introducing unnecessary difficulties. In Section 3 we point out that $\mathcal{T}_n$ is but a “visualization” of a more abstract geometric object, whose theory was still young at the epoch of von Dyck, namely a *colored graph* (Lemma 3.1). This viewpoint allows one to regard $D(n, n, n)$ as a group of colored graph automorphisms and to free oneself from the necessity to “realize” it over $\mathbb{S}$.

Once the definition of $D(n, n, n)$ has been detached from the notion of a tiling, it is easier to represent all its factors as group of automorphisms of a suitable quotient graph. This eliminates all problems coming from singularities which may be developed when $\mathbb{S}$ is factored by a discrete group of isometries.

Interestingly enough, such an interpretation of $D(n, n, n)$ as a group of colored graph automorphisms fits into a larger picture where *any* group D , equipped with a minimal additional structure, can be made act on a colored graph $T(D)$. The construction is discussed in Section 2, where we introduce (Definition 2.1) the key notion of a Borel-free set of generators of D , whose presence is indispensable to obtain the corresponding colored graph $T(D)$ (Proposition 2.1). This idea is rather old, and can be traced back to a 1962 paper by Tits [23] on the so-called *coset geometries* (see, e.g., [9, 4] and references therein). We prefer to regard a Tits geometry of rank m as a colored graph with m colors, since, to

¹The authors found this in [24, 8], but possibly it can be found somewhere else.

our opinion, it is more evident that a tiling determines a graph rather than a Tits geometry.

Moreover, the notion of a *chamber*, known in the context of Tits geometry (see [18, 21] and references therein), has a natural counterpart in the theory of colored graphs, namely that of a *clique*. Details are discussed in the first Section 1. By the definition of a Borel-free set of generators, the group D acts freely and transitively on the set of cliques $\mathfrak{C}(T(D))$ of $T(D)$, thus allowing to identify D with such a set (Lemma 1.2). Moreover, any factor of D can be identified with a quotient set of $\mathfrak{C}(T(D))$ (Corollary 2.1.1).

In Section 4 we show that, in the case of $D = D(n, n, n)$, the graph $T(D)$ is precisely the one obtained from the $2n$ -gonal tiling $\mathcal{P}_n$ of $\mathbb{S}$ determined by the triangular tiling $\mathcal{T}_n$ in the classical definition of a von Dyck group (Corollary 4.1.2). This leads to suspect that the very definition of a von Dyck group can be generalized in such a context, providing a natural “covering group” for $B(m, n)$ for $m > 2$. We discuss such a possibility in the final Subsection 7.1. To the authors knowledge, the fact that $T(D)$ is the planar graph determined by $\mathcal{P}_n$, has never been observed before, though a very similar result can be found in [24], presented a proprietary language.

Section 4 contains the main result of the paper, dubbed “cliques enumeration algorithm” (Corollary 4.2.1), which allows to recursively enumerate the cliques of the graph $T(D)$ and, hence, the elements of $D(n, n, n)$. In one form or another, any existing algorithm to check the finiteness of Burnside groups must implement a mechanism to enumerate words, and, due to the presence of relations, the lexicographic way is not necessarily the cheapest one (see [13] and references therein). The geometric understanding of $D(n, n, n)$ in terms of colored graphs allows to “avoid” relations and to produce exactly once each element of the group. To the authors opinion, this may eventually lead to effective techniques to attack the Burnside problem, and final Section 7 sketches the basic idea of a finiteness algorithm (Corollary 7.1.1).

The identification of $B(2, n)$ with a quotient set of $\mathfrak{C}(T(D))$ allows to exploit the cliques enumeration algorithm to check the finiteness of $B(2, n)$, since the elements of K_n can be enumerated as well (Corollary 6.0.2). Details are discussed in Section 6, where a geometric evidence of $B(2, 3)$ is provided (Subsection 6.1). We stress that the proposed enumeration of K_n is not injective, but this may be improved in future research.

Even if the computational cost of the proposed approach is still to be determined, especially in comparison with the existing techniques, it has the “aesthetic” merit of unifying the problems of finiteness of $B(2, n)$ for all values of n , in sharp contrast with the methods applied so far to concrete cases. Two classical proofs are reviewed in Subsections 5.1 and 5.2 of Section 5, which

collects some necessary facts about free Burnside groups and the Burnside problem.

Notation and conventions. All groups are assumed to be acting from the left. If the group D acts on the set S , then $\frac{S}{D}$ denotes the orbit space, and S is called a D -set. When needed, we write $d \cdot s$ for the result of the action of $d \in D$ on $s \in S$. Symbol $\frac{D}{K}$ denotes the quotient group with respect to a normal subgroup $K \trianglelefteq D$, or the space of left-cosets, when $K \leq D$ is an arbitrary subgroup.

A subset $Q \subseteq S$ such that each D -orbit intersects S exactly once is called a *fundamental set* of K (in S); if S is a surface, and D acts by isometries, we call a *fundamental domain* of K the closure of any fundamental set.

By a *tiling* $\mathcal{T}$ of a constant-curvature surface $\mathbb{S}$ we shall mean a family of regular closed and convex (geodesic) polygons in $\mathbb{S}$ such that the elements of $\mathcal{T}$ are pairwise isometrically equivalent and non-overlapping, and cover the whole $\mathbb{S}$ (see [11, 2]). Two polygons $P, Q \in \mathcal{T}$ are *non-overlapping* if the intersection $P \cap Q$ is either empty, or coincides with a vertex, or a whole edge. Any subset $\mathcal{T}_0 \subseteq \mathcal{T}$ will be referred to as a *sub-tiling* of $\mathcal{T}$. Notice that $\mathcal{T}_0$ fails, in general, to be a tiling of $\mathbb{S}$; it is a tiling of the closed subset $\mathcal{T}_0^\cup := \cup_{P \in \mathcal{T}_0} P \subseteq \mathbb{S}$. Observe that a sub-tiling is made of the same tiles of the parent tiling: not smaller, but less in quantity. All tilings considered in this paper will be made of even-sided polygons, except for an auxiliary tiling made of regular triangles (which is “inscribed” into all the polygonal ones). By the *boundary* of a sub-tiling $\mathcal{T}_0$ we mean the boundary of the closed subset $\mathcal{T}_0^\cup$, not to be mistaken with the graph of a tiling defined below.

Concerning graph terminology, we follow that of modern textbooks (see, e.g., [6, 10]): a graph is denoted by the symbol $\Gamma \equiv (V_\Gamma, E_\Gamma)$. We tacitly discard all the loops from the pseudo-graph associated to a symmetric and reflective binary relation, in such a way that it can be regarded as an ordinary graph. The notion of quotient graph is somehow rare in literature (see, e.g., [19]), and we quickly review it below; notions of graph (homo)morphisms and automorphisms can be found, e.g., in [6].

By the *graph* $\partial\mathcal{T}$ of a tiling $\mathcal{T}$, we mean the graph whose vertices (resp., edges) are the vertices (resp., edges) of the polygons in $\mathcal{T}$, hoping that the symbol “ ∂ ” will not mislead the reader.

We stress that our construction of a graph associated to a group is different from the usual Cayley graph.

Symbol n always denotes an integer ≥ 2 .

1. COLORED GRAPHS AND THEIR MORPHISMS

We review the notion of a *colored graph* and some related mathematical gadgets (see, e.g., [6]) stressing that some details, especially the definition of a *clique* may vary from other sources.

Definition 1.1. Let $\Gamma \equiv (V_\Gamma, E_\Gamma)$ be a graph, C a nonempty finite set, $\chi : V_\Gamma \rightarrow C$ a surjective map such that

$$(1.1) \quad \{v_1, v_2\} \in E_\Gamma \Rightarrow \chi(v_1) \neq \chi(v_2).$$

The pair (Γ, χ) is a *colored graph*. Elements of C are called *colors*: $\chi(v)$ is the *color* of the vertex $v \in V_\Gamma$, and χ is the *coloring function*. The *rank* of (Γ, χ) is the number of colors, i.e., $|C|$.

Roughly speaking, a colored graph is a graph where a color has been assigned to each vertex, in such a way that two vertices of the same color never form an edge. Plainly a given graph Γ can be colored in different ways: the minimum rank of all colored graphs of the form (Γ, χ) is the *chromatic number* of Γ . In order to formalize these “different ways” to color a graph, it is useful to introduce morphisms. We warn the reader that a pair (Γ, χ) will be identified with Γ in non ambiguous contexts.

Let now (Γ_i, χ_i) be a colored graph, $i = 1, 2$, and $F : \Gamma_1 \rightarrow \Gamma_2$ be a graph morphism. Roughly speaking, F is a morphism of colored graphs if it preserves the coloring.

Definition 1.2. F is a *morphism of colored graphs* if a map $f : C_1 \rightarrow C_2$ exists, such that the diagram

$$(1.2) \quad \begin{array}{ccc} \Gamma_1 & \xrightarrow{F} & \Gamma_2 \\ \downarrow \chi_1 & & \downarrow \chi_2 \\ C_1 & \xrightarrow{f} & C_2 \end{array}$$

is commutative. If f is a bijection, the morphism F is called *regular*. A regular morphism of the form (id_Γ, f) is called a *color shuffling*.

Together with morphisms, colored graphs form the *category of colored graphs*, denoted by **C-Gra**. For any $\Gamma \in \mathbf{C-Gra}$, the monoid $\text{Mor}(\Gamma, \Gamma)$ contains the distinguished subgroup $\text{Aut}(\Gamma)$ made of regular morphisms of the form (F, id_C) , where $F : \Gamma \rightarrow \Gamma$ is a graph automorphism.

In contexts where properties of colored graphs do not depend on the actual choice of colors, it is convenient to work with a category where the colorings are defined up to permutations. This is achieved as follows. First, take the sub-class of colored graphs of a fixed rank m : together with regular morphisms, it forms a (not full) sub-category of **C-Gra**. Then, in the so-obtained category, identify two morphisms if they differ by a color shuffling. The result is a new category, which we call the *category of m -colors graphs* and denote by **C-Gra_m**: an its object may be thought of as a colored graph Γ of rank m whose vertices are colored only up to permutations.

A sub-graph $\Gamma_0 \subset \Gamma$, can be understood as a sub-object in $\mathbf{C}\text{-Gra}_m$ only if the canonical inclusion $\iota : \Gamma_0 \rightarrow \Gamma$ gives rise to a morphism: in particular, Γ_0 must possess all colors of Γ . For example, if Γ_0 is a *complete sub-graph* (i.e., all its vertices are connected by an edge), it cannot be considered as a sub-object unless it has exactly m vertices.

Definition 1.3. A complete sub-graph $\Gamma_0 \subset \Gamma$ with m vertices is called a *clique*.

Notice that some authors (e.g., [16]) use the term “clique” to denote a complete sub-graph: if such a viewpoint was adopted then one should speak of a “clique with m vertices” instead, causing a pointless overload of the notation.

Hence, in the category $\mathbf{C}\text{-Gra}_m$, a clique Γ_0 is a sub-object of Γ . It is worth observing that any morphism $F : \Gamma \rightarrow \Gamma'$, being regular, is clique-preserving (i.e., $F(\Gamma_0)$ is a clique in Γ') and as such it can be restricted to the sub-object Γ_0 .

Denote by $\mathfrak{C}(\Gamma)$ the (possibly empty) set of all cliques of Γ .

Remark 1. Let $\mathfrak{C}(\Gamma) \neq \emptyset$. For similar reasons as above, any $F \in \text{Aut}(\Gamma)$ is clique-preserving (F is a graph automorphism) and as such it induces a bijection $\mathfrak{C}(F)$ on $\mathfrak{C}(\Gamma)$. Correspondence $\mathfrak{C} : F \mapsto \mathfrak{C}(F)$ is a group homomorphism between $\text{Aut}(\Gamma)$ and the group of permutations of $\mathfrak{C}(\Gamma)$.

Definition 1.4. $\text{Aut}(\Gamma)$ is the *symmetry group* of Γ ; its elements are called *transformations* of Γ . A group homomorphism $i : D \rightarrow \text{Aut}(\Gamma)$ is called an *action* of D on Γ , which is *clique-transitive* (resp., *clique-free*) if $\mathfrak{C}(\Gamma) \neq \emptyset$ and the action $\mathfrak{C} \circ i$ of D on $\mathfrak{C}(\Gamma)$ is transitive (resp., free).

If D acts on (Γ, χ) then the *quotient colored graph* $(\frac{\Gamma}{D}, \tilde{\chi})$ is defined as follows:
 $V_{\frac{\Gamma}{D}} := \frac{V_{\Gamma}}{D}$,

$$(1.3) \quad (Dv_1, Dv_2) \in E_{\frac{\Gamma}{D}} \Leftrightarrow \exists d_1, d_2 \in D \mid (d_1 v_1, d_2 v_2) \in E_{\Gamma},$$

and $\tilde{\chi}(Dv) := \chi(v)$.

Lemma 1.1. *The canonical projection $\pi : V_{\Gamma} \rightarrow V_{\frac{\Gamma}{D}}$ is a morphism in $\mathbf{C}\text{-Gra}$. Moreover, if $\tilde{v}_1$ and $\tilde{v}_2$ are connected by an edge in $\frac{\Gamma}{D}$, then any $v_1 \in \pi^{-1}(\tilde{v}_1)$ is connected by an edge with an element of $\pi^{-1}(\tilde{v}_2)$.*

Proof. By (1.3) π is manifestly edge-preserving and color-preserving: as such, it defines a morphism in $\mathbf{C}\text{-Gra}$ (see Definition 1.2). Let now $(\tilde{v}_1, \tilde{v}_2) \in E_{\frac{\Gamma}{D}}$. Then $v'_1 \in \pi^{-1}(\tilde{v}_1)$ and $v_2 \in \pi^{-1}(\tilde{v}_2)$ exist, such that $(v'_1, v_2) \in E_{\Gamma}$. But D acts on Γ by graph automorphisms, and transitively on $\pi^{-1}(\tilde{v}_1)$: so, $v_1 = dv'_1$, with $d \in D$, and hence $(v_1, dv_2) = d(v'_1, v_2) \in E_{\Gamma}$. $\square$

Remark 2. Observe that π induces a map $\pi_* : \mathfrak{C}(\Gamma) \rightarrow \mathfrak{C}(\frac{\Gamma}{D})$ which is, in general, not surjective. Let $\tilde{c} \in \mathfrak{C}(\frac{\Gamma}{D})$ be a clique of $\frac{\Gamma}{D}$, and $V_{\tilde{c}} = \{\tilde{v}_1, \dots, \tilde{v}_m\}$

be the set of its vertices. Then, by Lemma 1.1, it is possible to show that $v_j \in \pi^{-1}(\tilde{v}_j)$ exists, for $j = 1, \dots, m$, such that v_1 forms an edge with $v_2, \dots, v_m$. So, $V_c := \{v_1, \dots, v_m\}$ does not, in general, determine a clique $c \in \mathfrak{C}(\Gamma)$ unless $m = 2$. In this paper we shall mainly be interested in the case $m = 2$, but future generalizations may require higher values of m : then, it will be needed to find extra conditions on the action of D guaranteeing surjectivity of π_* .

Observe that a clique-transitive action does not need to be edge-transitive and it is *never* vertex-transitive, since vertices of different colors cannot be mapped one into another by an element of $\text{Aut}(\Gamma)$. Hence, if D acts clique-transitively, the quotient graph $\frac{\Gamma}{D}$ reduces to a single clique, which, in general, contain more than one edge (and, hence, more than two vertices!). Lemma 1.2 below finalizes this section with a list of useful results.

Lemma 1.2. *Let D be a group acting clique-freely and clique-transitively on Γ , and $K \trianglelefteq D$ be a normal subgroup of D . Then,*

- (1) *D is D -equivariantly identified with the set $\mathfrak{C}(\Gamma)$ of cliques of Γ ,*
- (2) *if a clique $c_0 \in \mathfrak{C}(\Gamma)$ is fixed then $\mathfrak{C}(\Gamma)$ has a group structure isomorphic to D , whose unity is c_0 .*

Moreover, if π_* is surjective, then

- (3) *the factor group $\frac{D}{K}$ acts clique-freely and clique-transitively on the quotient graph $\frac{\Gamma}{K}$,*
- (4) *$\frac{D}{K}$ is $\frac{D}{K}$ -equivariantly identified with the quotient set $\frac{\mathfrak{C}(\Gamma)}{K} = \mathfrak{C}(\frac{\Gamma}{K})$.*

Proof. By Definition 1.4 D acts freely and transitively on the set $\mathfrak{C}(\Gamma)$: hence (1) and (2) are valid. Moreover, since $\frac{D}{K}$ acts freely and transitively on $\frac{\mathfrak{C}(\Gamma)}{K}$, the group $\frac{D}{K}$ is $\frac{D}{K}$ -equivariantly identified with the set $\frac{\mathfrak{C}(\Gamma)}{K}$. In other words, in order to prove (4), it is enough to prove (3).

To this end, let $\tilde{c} \in \mathfrak{C}(\frac{\Gamma}{K})$ be a clique of $\frac{\Gamma}{K}$, and write it as $\tilde{c} = \pi_*(c)$. Then the desired clique-free and clique-transitive action of $\frac{D}{K}$ on $\frac{\Gamma}{K}$ is defined by

$$(1.4) \quad \begin{aligned} \frac{D}{K} \times V_{\frac{\Gamma}{K}} &\longrightarrow V_{\frac{\Gamma}{K}}, \\ (dK, \pi(v)) &\longmapsto \pi(dv). \end{aligned}$$

First, we check correctness of (1.4): if $(d'K, \pi(v')) = (dK, \pi(v))$ then $d' = dk$ and $v' = k_1v$, with $k, k_1 \in K$. So, $\pi(d'v') = \pi(dkk_1v) = \pi(k_2dv) = \pi(dv)$, where $k_2 \in K$ is such that $(kk_1)^{d^{-1}} = k_2$.

Observe now that (1.4) is edge-preserving. Indeed, by Lemma 1.1, if $(\pi(v_1), \pi(v_2)) \in E_{\frac{\Gamma}{K}}$, then it is possible to assume that $(v_1, v_2) \in E_\Gamma$. But K acts by graph automorphisms, so that $d(v_1, v_2) = (dv_1, dv_2) \in E_\Gamma$ as well, and then $(\pi(dv_1), \pi(dv_2)) \in E_{\frac{\Gamma}{K}}$.

Let us verify clique-transitivity. Given two cliques $\tilde{c}, \tilde{c}_1 \in \mathfrak{C}(\frac{\Gamma}{K})$, write them as $\tilde{c} = \pi_*(c)$, $\tilde{c}_1 = \pi_*(c_1)$, for $c, c_1 \in \mathfrak{C}(\Gamma)$. Then, by clique-transitivity of D -action, $c_1 = dc$, for some d , and hence $dK \cdot \tilde{c} = \tilde{c}_1$.

Finally, if $dK \cdot \tilde{c} = \tilde{c}$, then $V_{\tilde{c}} = \{\pi(dv_1), \dots, \pi(dv_m)\} = \{\pi(v_1), \dots, \pi(v_m)\}$, i.e., $\pi(dv_j) = \pi(v_j)$ for all $j = 1, \dots, m$, since v_j and dv_j have the same color: hence $d \in K$. $\square$

2. THE COLORED GRAPH OF A GROUP WITH A BOREL-FREE SET OF GENERATORS

A group D is said to be the direct (inner) product of its normal subgroups H and K , if $D = \langle H, K \rangle$ and $H \cap K = 1$. We turn our attention to those generator sets $S \subseteq D$ which, in a sense, allow to express D as the “direct product” of the cyclic groups generated by the elements of S .

Definition 2.1. S is a *Borel-free* set of generators if $\bigcap_{s \in S} \langle s \rangle = 1$, and $\langle s \rangle \neq 1$ for all $s \in S$.

Observe that, in the case $S = \{x, y\}$ and $\langle x \rangle, \langle y \rangle \trianglelefteq D$, Definition 2.1 states precisely that D is the direct product of $\langle x \rangle$ and $\langle y \rangle$.

Example 1. The group of quaternions Q does not admit any Borel-free set of generators, since any nontrivial subgroup of Q contains the cyclic subgroup $Z(Q) = \{\pm 1\}$.

Example 2. Let $D := \langle a, b \mid a^{p^2} = b^p = 1, a^b = a^{1+p} \rangle$ the extra-special group of exponent p^2 (see [20]) and $S := \{a, ab\}$, with p odd prime. Then S generates D (since $b = a^{-1}ab$) but it is not Borel-free because² $1 \neq a^p = (ab)^p$, and hence $\langle a \rangle \cap \langle ab \rangle \neq 1$. It follows that not all set of generators of rank 2 are necessarily Borel-free.

A group D with a Borel-free set of generators S of rank m allows to define a colored graph $T(D)$, of same rank m , in the following way:

- (1) the vertices of $T(D)$ are the left-cosets of the subgroups $\langle s \rangle$, for all $s \in S$;
- (2) two vertices are connected by an edge if the corresponding cosets have nonempty intersection;
- (3) the set of colors is S ;
- (4) the coloring function $\chi : V_{T(D)} \rightarrow S$ is the map unambiguously defined by

$$\chi(d\langle s \rangle) := s, \quad d \in D.$$

²This equality can be easily obtained from the identity $(ab)^n = a^n b^n [b, a]^{\binom{n}{2}}$ which in turn is due to the fact that $[a, b] = a^p$ is central.

Proposition 2.1. *Let S be a Borel-free set of generators of D , and define $T(D)$ and χ as above. Then,*

- (1) *$(T(D), \chi)$ is a colored graph of rank m ;*
- (2) *D acts clique-transitively and clique-freely on $(T(D), \chi)$;*
- (3) *any element $d \in D$ acts on cliques by a sequence of rotations.*

Proof. As the disjoint union of the left D -sets $\frac{D}{\langle s \rangle}$, $s \in S$, the set of vertices $V_{T(D)}$ inherits a left D -structure. Observe also that D acts transitively on each subset of vertices $\frac{D}{\langle s \rangle}$, i.e., it is color-preserving, and that the sub-graph $c_0 := \{\langle s \rangle \mid s \in S\}$ is a clique; indeed, any two its vertices are connected by an edge, and any vertex outside c_0 is of the form $d \cdot \langle s \rangle$, for some $s \in S$ and $d \in D \setminus \langle s \rangle$ and, as such, it cannot be connected by an edge to $\langle s \rangle$, since they have the same color s . Finally, the action of D sends edges to edges, since two nontrivially intersecting left cosets are mapped into nontrivially intersecting left cosets, i.e., $D \leq \text{Aut}(T(D))$.

Prove that any clique c is of the form $d \cdot c_0$. Take a vertex $d \cdot \langle s \rangle \in c$ and observe that $d^{-1} \cdot c$ is a clique containing $\langle s \rangle$ (Remark 1). By Definition 1.3 it must coincide with c_0 . Hence, D acts clique-transitively. Finally, $d \in D$ stabilizes c_0 if and only if it stabilizes all its vertices, i.e., it belongs to $\bigcap_{s \in S} \langle s \rangle = 1$.

It remains to be observed that $d = s$ acts on c_0 by a rotation around its vertex $\langle s \rangle$. Let now $d = \underline{d}s$, and suppose that $\underline{d}$ acts on c_0 by a sequence of rotations: then, $d \cdot c_0 = (\underline{d}s\underline{d}^{-1}) \cdot (\underline{d} \cdot c_0)$ is obtained from c_0 by the *same* sequence of rotations, followed by a rotation around the vertex $\underline{d} \cdot \langle s \rangle$, since $\underline{d}\langle s \rangle\underline{d}^{-1}$ is precisely the stabilizer of $\underline{d} \cdot \langle s \rangle$. $\square$

Observe that (1) is just a paraphrase of the fact that $T(D)$ is the colored graph associated to a coset geometry (in the sense of Tits–Buekenhausen [23, 4] where *chambers* play the role of cliques) of rank m , (2) follows from the triviality of the intersection of the $\langle s \rangle$'s, and (3) is a consequence of the fact that $D = \langle S \rangle$.

It is worth stressing that “rotation” refers to a transformation of Γ with a fixed point (vertex) v ; hence, it does need to be of finite order: geometrically, this means that there may be infinitely many edges with one vertex equal to v .

Correspondence $(D, S) \mapsto (T(D), \chi)$ has a manifest functor character. For example, let $\pi : D \rightarrow B$ be an epimorphism and S (resp., $\underline{S}$) a Borel-free generator set of D (resp., B) such that $\pi(S) = \underline{S}$ and $\pi|_S : S \rightarrow \underline{S}$ is a bijection. Then the following result is a direct consequence of Lemma 1.2 (4) (see also Remark 2).

Observe that if $m = 2$ then D is the set of the oriented edges of a bipartite graph.

Corollary 2.1.1. *If $m = 2$ then the colored graph $T(B)$ coincides with the quotient graph $\frac{T(D)}{\ker \pi}$ and hence B is B -equivariantly identified with the quotient set $\frac{\mathcal{C}(T(D))}{\ker \pi}$.*

The next step is to show that for some groups D with two generators $S = \{x, y\}$ the bipartite graph $T(D)$ actually belongs to a richer geometrical object, namely a surface, thus giving a more concrete meaning to the notion of “rotation”.

By Corollary 2.1.1 if B is “covered” by a larger group D , whose colored graph $T(D)$ takes a particular simple form like, e.g., the boundary of a regular tiling, then B can be identified with a quotient of the set of cliques of $T(D)$. In the next sections the role of “covering group” D will be played by a von Dyck group while B is the two-generators free Burnside group.

3. $2n$ -GONAL TILING OF CONSTANT-CURVATURE SURFACES AND THEIR 2-COLORS GRAPHS

From now on, symbol $\mathbb{S}$ will denote a constant-curvature surface, i.e., $\mathbb{S} = S^2, \mathbb{R}^2$, or $\mathbb{H}$, and $\mathcal{T}_n$ an its tiling by regular triangles of angle $\frac{\pi}{n}$ (see, e.g., [2]). Obviously $\mathcal{T}_2$ exist only on S^2 , $\mathcal{T}_3$ only on $\mathbb{R}^2$, and all the $\mathcal{T}_n$'s, with $n > 3$, exists only on $\mathbb{H}$. We suppose also that there is a distinguished triangle $\Delta_n \in \mathcal{T}_n$, called *basic*, and a distinguished vertex $O \in \Delta_n$ called *center* (of the tiling). Due to homogeneity, such choices are by no means restrictive.

The *basic $2n$ -gon* P_n of a tiling $\mathcal{T}_n$ is the union of all triangles obtained by rotating the basic triangle around the center of the tiling.

Lemma 3.1. *A unique 3-colors complete graph $\partial\mathcal{T}_n \in \mathbf{C}\text{-}\mathbf{Gra}_3$ exists whose vertices (resp., edges) are the vertices (resp., edges) of $\mathcal{T}_n$.*

Proof. The existence of the graph is obvious. The coloring function $\chi : V_{\partial\mathcal{T}_n} \rightarrow \{\bullet, \circ, \triangle\}$ can be arbitrarily defined on the vertices of the basic triangle. Suppose now that χ has been defined on a subgraph $\Gamma' \subset \partial\mathcal{T}_n$ and let $\Delta \in \mathcal{T}_n$ be a triangle with two vertices in Γ' : then there is a unique way to define χ on the third vertex. Since $\partial\mathcal{T}_n$ is an object in $\mathbf{C}\text{-}\mathbf{Gra}_3$, χ can be defined up to chromatic permutations. $\square$

Starting from the basic $2n$ -gon P_n define a unique $2n$ -gonal tiling $\mathcal{P}_n$ of $\mathbb{S}$. We call P_n the *basic $2n$ -gon* of $\mathcal{P}_n$ and O its center. Observe that O is not a vertex of P_n . Define the unique 2-colors (i.e., bipartite) subgraph $\partial\mathcal{P}_n \subseteq \partial\mathcal{T}_n$, by taking only the vertices and edges belonging to $\mathcal{P}_n$. Observe that only $\mathcal{T}_2$ is finite, consisting of 8 triangles; hence $\mathcal{P}_2$ consists of 2 squares, namely the upper and lower hemisphere of S^2 .

Therefore, on a constant-curvature surface $\mathbb{S}$ there is a 2-colors graph $\partial\mathcal{P}_n$, whose edges are geodesic segments of a fixed length, which originated as the

boundary of a regular $2n$ -gonal tiling $\mathcal{P}_n$, whose tiles, in turn, were obtained by the (regular) triangle tiles of a regular triangular tiling $\mathcal{T}_n$. This last observation is a compulsory step to make von Dyck group act on the 2-colors graph $\partial\mathcal{P}_n$. In Section 4 below we show that such an action is clique-free and clique-transitive, allowing to use the results from Sections 1 and 2.

Observe that in a 2-colors graph a clique is nothing but an edge: nevertheless, clique-transitivity must not be mistaken for edge-transitivity since a clique-transitive action can never switch the colors of the ends of an edge. It is also worth mentioning that $\partial\mathcal{P}_n$ is made of *loops* consisting of $2n$ vertices of alternating colors (they correspond exactly to the boundaries of the tiles of $\mathcal{P}_n$): a transformation of $\partial\mathcal{P}_n$ cannot be realized as a rotation of an angle $\frac{\pi}{n}$, since it would switch vertex colors.

4. TRIANGLE AND VON DYCK GROUPS

In the previous sections we proposed an abstract construction to associate a 2-colors graph with a 2-generators group. We show now that in the case of special discrete subgroups of isometries of $\mathbb{S}$ such a graph can be realized by means of a polygonal tiling of $\mathbb{S}$ (Corollary 4.1.2).

Definition 4.1. Let a, b, c positive integers. The von Dyck group (see [25] and references therein) is

$$(4.1) \quad D(a, b, c) := \langle x, y \mid x^a = y^b = (xy)^c = 1 \rangle.$$

Groups (4.1) were introduced by Walther von Dyck at the end of eighteen century.

Theorem 4.1 ([2]). Let $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} > 1$ (resp. $= 1, < 1$). Then $D(a, b, c)$ is the group of orientation-preserving transformations of a tiling of $\mathbb{S} = S^2$ (resp., $\mathbb{R}^2, \mathbb{H}$) generated by the rotations around the vertices of a basic triangle.

Corollary 4.1.1 ([2]). Let $n > 3$. Then $D(n, n, n)$ is a Fuchsian group, i.e., a discrete subgroup of orientation-preserving isometries of $\mathbb{H}$.

Recall that, if orientation-reversing transformations are taken into account, one obtains a larger group, the so-called *triangle group* (see, e.g., [1] and references therein) containing $D(a, b, c)$ as a subgroup of index 2. Observe that the triangle group acts transitively on $\mathcal{T}_n$, unlike $D(n, n, n)$, which cannot move a triangle into an adjacent one.

Corollary 4.1.2. The von Dyck group $D(n, n, n)$ acts clique-transitively and clique-freely on the 2-colors graph $\partial\mathcal{P}_n$. Moreover, $\partial\mathcal{P}_n$ is canonically and $D(n, n, n)$ -equivariantly identified with $T(D(n, n, n), \{x, y\})$.

Proof. $D(n, n, n)$ acts on the 3-colors graph $\partial\mathcal{T}_n$ (see Lemma 3.1) and the stabilizer of the basic triangle is trivial, i.e., the action is clique-free. Even if the




FIGURE 1. The same element d of the von Dyck group $D(n, n, n)$ acts both on the oriented triangles of the regular tiling $\partial(\mathcal{T}_n)$, and on cliques of a 2-colors graph $T(D(n, n, n), \{x, y\})$.

action is not clique-transitive, two cliques of the same *orientation* are always moved one to another by an element of $D(n, n, n)$ (Theorem 4.1). It remains to observe that a clique of $\partial\mathcal{P}_n$ determines a unique oriented clique of $\partial\mathcal{T}_n$: it is enough to choose a fixed sequence of colors, say red-green-blue and, to any clique of $\partial\mathcal{P}_n$, i.e., an edge with red-green ends, associate the unique (out of the two adjacent triangles) clique of $\partial\mathcal{T}_n$ whose blue vertex makes the resulting sequence red-green-blue run counterclockwise. Figure 1 shows two triangles facing the same clique (A, B) : only the colored one has positive orientation.

By Proposition 2.1 any clique c of $T(D(n, n, n), \{x, y\})$ is uniquely written as $c = (d\langle x \rangle, d\langle y \rangle)$, with $d \in D(n, n, n)$. Let (A, B, C) the oriented clique of $\partial\mathcal{T}_n$ determined by $A = d\langle x \rangle$ and $B = d\langle y \rangle$: hence, (A, B, C) can be uniquely written as $d'(A_0, B_0, C_0)$, where $A_0 = \langle x \rangle$ and $B_0 = \langle y \rangle$ (see Figure 1). It follows that $d' = d$. But the oriented clique (A, B, C) is uniquely determined by the clique (A, B) of $\partial\mathcal{P}_n$: correspondence $c = (d\langle x \rangle, d\langle y \rangle) \leftrightarrow (A, B) = (dA_0, dB_0)$ is manifestly one-to-one and $D(n, n, n)$ -equivariant.

□

By Lemma 1.2 $D(n, n, n)$ can be identified with the set of cliques $\mathfrak{C}(\partial\mathcal{P}_n)$ and all its factors with a quotient of $\mathfrak{C}(\partial\mathcal{P}_n)$. In the next section we show that among the factors of $D(n, n, n)$ there are the free Burnside groups with two generators. The generalization of this picture to the cases with more generators is a challenging task. It is worth stressing that only for $n = 2$ the set of cliques $\mathfrak{C}(\partial\mathcal{P}_n)$ is finite.

From now on the generator x (resp. the generator y , the product $r := xy$) will be identified with the $\frac{2\pi}{n}$ rotation of $\mathcal{P}_n$ around the vertex H (resp., the vertex K , the center O), where (O, H, K) is a fixed orientation of the basic triangle (Theorem 4.1). We call $c_0 = (H, K)$ the *basic clique* and we identify any element $w \in D(n, n, n)$ with the clique $wc_0 = (wH, wK)$.

By Proposition 2.1 $D(n, n, n)$ acts on $\partial\mathcal{P}_n$ by a sequence of rotations.

Remark 3. $D(n, n, n)$ acts transitively on polygons of $\mathcal{P}_n$ since these are “minimal loops”, but the stabilizers are the cyclic subgroups of order n conjugate to $\langle r \rangle$.

Remark 4. When a loop Γ is oriented any its clique $c \in \mathfrak{C}(\Gamma)$ inherits an orientation: we say that c is *positively* (resp., *negatively*) *oriented* if, running accordingly to the positive orientation of Γ , the x -colored vertex comes after the y -colored one (resp., before). Observe that any loop must be composed of an even number of cliques: of these, half is positively oriented, and the other half is negatively oriented.

Remark 5. Denote by $\lfloor \frac{i}{2} \rfloor$ the integer part of $\frac{i}{2}$ and by $[i]_{2\mathbb{Z}}$ the parity of i . Then the map

$$i \in \{0, 1, \dots, 2n-1\} \mapsto d(i) := r^{\lfloor \frac{i}{2} \rfloor} x^{[i]_{2\mathbb{Z}}} \cdot c_0 \in \mathfrak{C}(\partial\mathcal{P}_n),$$

is a bijection.

Proposition 4.2 (Polygonal enlargement). *Let $\mathcal{Q} \subseteq \mathcal{P}_n$ be a sub-tiling such that its boundary is a simple (i.e., not self-intersecting) loop $\Gamma \subseteq \partial\mathcal{P}_n$ consisting of N cliques. Define $\tilde{\mathcal{Q}}$ as the larger sub-tiling obtained from $\mathcal{Q}$ by adding all the $2n$ -gons which intersect it and call $\tilde{\Gamma}$ its boundary. Assume both Γ and $\tilde{\Gamma}$ to be oriented clockwise. Then*

- (1) *for any positively oriented clique $w_0 \in \mathfrak{C}(\Gamma)$ there exist unique integers $i_1, \dots, i_N \in \{1, \dots, n-1\}$ such that the sequence*

$$(4.2) \quad w_0, w_0 x^{i_1}, w_0 x^{i_1} y^{i_2}, \dots, w_0 x^{i_1} y^{i_2} \dots x^{i_{N-1}} y^{i_N} = w_0$$

run clockwise through all the cliques of Γ ;

- (2) *the number of $2n$ -gons contained in $\tilde{\mathcal{Q}} \setminus \mathcal{Q}$ is*

$$(4.3) \quad N(n-1) - \sum_{k=1}^N i_k;$$

- (3) *by replacing in (4.2) each positively (resp., negatively) oriented clique $w = w_0 \dots y^{i_{k-1}}$ (resp., $v = w_0 \dots x^{i_k}$) according to the following rules*

$$(4.4) \quad w \leftrightarrow \underbrace{(wx^{n-2}d(2), \dots, wx^{n-2}d(2n-1), \dots, wx^{i_k}d(2), \dots, wx^{i_k}d(2n-2))}_{\text{not present for } n-2=i_k},$$

$$(4.5) \quad v \leftrightarrow \underbrace{(vy^{n-1}d(1), \dots, vy^{n-1}d(2n-2), \dots, vy^{i_{k+1}+1}d(1), \dots, vy^{i_{k+1}+1}d(2n-3))}_{\text{not present for } n-1=i_{k+1}+1},$$

one obtains the (positively oriented) sequence of cliques of $\tilde{\Gamma}$.

Proof. The cliques of Γ are of alternating orientation (Remark 4) and any of them can be obtained by another by a sequence of rotations (Proposition 2.1 (3)). In particular, the clique coming next to w (which is, as such, negatively oriented) is obtained by a rotation around the x -colored vertex of w : hence it is uniquely written as wx^{i_k} (see the proof of point (3) of Proposition 2.1). By iteration one obtains the sequence (4.2), proving (1).

Figure 2 shows the n cliques pivoted at the x -color vertex of the (positively oriented) clique w . Starting from w itself and running counterclockwise one meets the internal cliques $wx, \dots, wx^{i_k-1}$ (highlighted in red) and then the next (negatively oriented) clique $v = wx^{i_k}$ of Γ . It is convenient (since they will form the loop $\tilde{\Gamma}$) to list clockwise the remaining $n - i_k - 1$ cliques, i.e., those outside $\mathcal{Q}$, namely

$$wx^{n-1}, wx^{n-2}, \dots, wx^{i_k+1}$$

(highlighted in blue).

Now the $2n$ -gon of $\mathcal{P}_n$ determined by the two consecutive cliques wx^{n-2} and wx^{n-1} is precisely $wx^{n-2}P_n$; continuing until the $2n$ -gon $wx^{i_k}P_n$, we obtain exactly $n - i_k - 1$ $2n$ -gons. The cliques $wx^{n-2} = wx^{n-2}d(0)$ and $wx^{n-1} = wx^{n-2}d(1)$ of $wx^{n-2}P_n$ will be inside $\tilde{\mathcal{Q}}$: hence, the remaining ones, i.e., $wx^{n-2}d(2), \dots, wx^{n-2}d(2n-1)$, are those which will contribute to $\tilde{\Gamma}$ (highlighted in green). Repeating this for $n-3, \dots, i_k+1$ explains the marked part of (4.4). The last $2n$ -gon, i.e., $wx^{i_k}P_n$ has three cliques inside $\tilde{\mathcal{Q}}$, namely $wx^{i_k}d(2n-1)$, $wx^{i_k}d(0) = wx^{i_k}$ (belonging to $\mathcal{Q}$) and $wx^{i_k}d(1)$: hence, the ones belonging to $\tilde{\Gamma}$ are $wx^{i_k}d(2), \dots, wx^{i_k}d(2n-2)$ (highlighted in purple) and this explains the remaining part of (4.4).

Let us pass to a negatively oriented clique v (for simplicity, $v = wx^{i_k}$). The (clockwise) list of the cliques outside $\mathcal{Q}$ obtained by rotating v around its y -vertex is

$$vy^{n-1}, vy^{n-2}, \dots, vy^{i_{k+1}+1}.$$

Again we obtain $n - i_k - 1$ $2n$ -gons. So, any vertex of Γ intersects exactly $n - i_k - 1$ $2n$ -gons lying outside $\mathcal{Q}$: hence, (4.3) holds true and (2) is proved.

Now observe that the $2n$ -gon determined the two consecutive cliques vy^{n-2} and vy^{n-1} is precisely $vy^{n-1}P_n$: indeed, the former is $vy^{n-1}d(2n-1)$ and the latter is $vy^{n-1}d(0)$. Hence, the remaining cliques $vy^{n-1}d(1), \dots, vy^{n-1}d(2n-2)$ are those which will contribute to $\tilde{\Gamma}$. Repeating this for $n-2, \dots, i_{k+1}+2$ one obtains the marked part of (4.5). The last $2n$ -gon, which is $vy^{i_{k+1}+1}P_n$, has

three cliques inside $\tilde{\mathcal{Q}}$: $vy^{i_{k+1}+1}d(2n-2)$, $vy^{i_{k+1}+1}d(2n-1) = vy^{i_{k+1}}$ (belonging to $\mathcal{Q}$) and $vy^{i_{k+1}+1}d(0)$. The remaining ones, $vy^{i_{k+1}+1}d(1), \dots, vy^{i_{k+1}+1}d(2n-3)$, explain the last part of (4.5) and are highlighted orange in Figure 2.

Observe that in the Euclidean case ($n = 3$) any hexagon intersecting $\mathcal{Q}$ must be adjacent to it: hence, the marked parts of (4.4) and (4.5) must be suppressed (compare Figures 4 and 6). $\square$

The meaning of the number i_k is that $2\pi \frac{i_k}{n}$ represents the internal angle of Γ at its k^{th} vertex (according to sequence (4.2)): on this concern, it is worth observing that the algorithm of Proposition 4.2 does not requires $\mathcal{Q}$ to be convex, i.e., large values of i_k are allowed (though not useful for our purposes). In the Euclidean case, $i_k = 2$ is already a “large” value: hence no hexagon at all will be added there by the algorithm; there are only three hexagons around any vertex, and if one is “inside” $\mathcal{Q}$ (i.e., $i_k = 1$) then the remaining two ones must be “outside” and both must have an edge in common with $\mathcal{Q}$ (whence the disappearance of the marked parts of (4.4) and (4.5), which would correspond to an hexagon having just one vertex in common with $\mathcal{Q}$).

Corollary 4.2.1 (Cliques enumeration algorithm). *There exists a recursively defined bijection $d : \mathbb{N}_0 \rightarrow D(n, n, n)$.*

Proof. For $n = 2$, the group $D(n, n, n)$ is finite (Theorem 4.1) so there is nothing to prove.

Let $n \geq 3$. Define d on $N_1 := \{0, 1, \dots, 2n-1\}$ as in Remark 5. Let $\widetilde{P}_n$ the enlargement of P_n constructed as in Proposition 4.2 and use the current definition of d to extend it to $N_1 \cup N_2$, where $N_2 \subseteq \mathbb{N}_0 \setminus N_1$ is in one-to-one correspondence with the set of cliques $\mathfrak{C}(\partial \widetilde{P}_n) \setminus \mathfrak{C}(\partial P_n)$. In order to fill up the whole $\mathbb{N}_0$ we choose N_2 in such a way that, together with N_1 , it forms an interval.

It remains to be noticed that the boundary of all successive enlargements of P_n are always regular, i.e., Proposition 4.2 can be used to recursively define the bijection d between $\mathbb{N}_0 = \cup_{j=1}^{\infty} N_j$ and $\mathfrak{C}(\partial \mathcal{P}_n) = \cup_{j=1}^{\infty} (\mathfrak{C}(\partial P_n^{(j+1)}) \setminus \mathfrak{C}(\partial P_n^{(j)}))$, where $P_n^{(j+1)} = \widetilde{P_n^{(j)}}$ and $P_n^{(1)} = P_n$. $\square$

Let now $n > 3$, and consider the j^{th} enlarged polygon $P_n^{(j)}$ defined in Corollary 4.2.1.

Corollary 4.2.2. *The elements $d(3)^j$ and $d(2n-3)^j$ belong to $\mathfrak{C}(\partial P_n^{(j)}) \setminus \mathfrak{C}(\partial P_n^{(j-1)})$ for all $j > 1$ and $[d(3)^n, d(2n-3)^n] \neq 1$.*

Proof. The first statement is a direct consequence of Proposition 4.2. To begin with, observe that the part of (4.4) which not underbraced, i.e.,




FIGURE 2. Polygonal enlargement procedure.

$wx^{i_k}d(2), \dots, wx^{i_k}d(2n-2)$, can be rewritten as

$$(4.6) \quad vd(2), vd(3), \dots, vd(2n-3), vd(2n-2),$$

with $v := wx^{i_k}$. Now we can continue the proof by induction on j . Obviously $d(3)^1$ and $d(2n-2)^1$ both belong to $\mathfrak{C}(\partial P_n)$. Let now $j > 1$ and assume that $d(3)^j$ and $d(2n-3)^j$ belong to $\mathfrak{C}(\partial P_n^{(j)}) \setminus \mathfrak{C}(\partial P_n^{(j-1)})$. In order to show that $d(3)^{j+1}$ and $d(2n-3)^{j+1}$ belong to $\mathfrak{C}(\partial P_n^{(j+1)}) \setminus \mathfrak{C}(\partial P_n^{(j)})$, it is enough to observe that

$$d(3)^{j+1} = \underbrace{d(3)^j}_{\in \mathfrak{C}(\partial P_n^{(j)}) \setminus \mathfrak{C}(\partial P_n^{(j-1)})} d(3) = xyx^2yx^2 \dots x^2yx d(3)$$

can be written as $vd(3)$ where $v := xyx^2yx^2 \dots x^2yx$, and as such it belongs to $\mathfrak{C}(\partial P_n^{(j+1)}) \setminus \mathfrak{C}(\partial P_n^{(j)})$, in view of (4.6). The same reasoning is valid for $d(2n-3)^{j+1}$.

The second statement is checked by explicitly computing the Möbius transformations associated with $d(3)^j = (xyx)^j$ and $d(2n-3)^j = ((xy)^{n-2}x)^j$. It is worth to notice that it would be impossible to solve the equation $[d(3)^n, d(2n-3)^n] = 1$ in a purely algebraic way, because of the unknown exponent n . On the other hand, the geometric interpretation allows us to apply the formula for the n^{th} (and $-n^{\text{th}}$) power of unimodular matrices based on Chebyshev polynomials of the second kind (see, e.g., [3]) to the elements uvu and $v^{-1}u^{-1}v^{-1} = (uv)^{n-2}u$, where the matrices

$$u = \begin{pmatrix} \cos \frac{\pi}{n} & -\sin \frac{\pi}{n} \\ \sin \frac{\pi}{n} & \cos \frac{\pi}{n} \end{pmatrix} \text{ and } v = \begin{pmatrix} \cos \frac{\pi}{n} & -\beta \sin \frac{\pi}{n} \\ (\sin \frac{\pi}{n})/\beta & \cos \frac{\pi}{n} \end{pmatrix}$$


represent a counterclockwise rotation of angle $2\pi/n$ about the vertices $x := i$ and $y := \beta i$, respectively. The real number β comes from the theory of regular polygons on hyperbolic plane: it is defined by the property that $[\beta i, i]$ is one of the sides of a regular $2n$ -gon; as such, β depends on the choice of the interior angle $\frac{2\pi}{n}$, and it can be found by solving the equation $\beta + \frac{1}{\beta} = \frac{2 \cos \frac{\pi}{n}}{1 - \cos \frac{\pi}{n}}$. Now the desired commutator can be computed in a matrix form³ and, to check that it is not identical, it is enough to consider, e.g., its $(1, 2)$ entry, say $a(n)$, and notice that the function a is strictly positive and growing for $n \geq 4$. □

Figure 3 shows how the cliques enumeration algorithm works for $n = 3$. The white hexagon is P_3 and its perimeter $\Gamma := \partial P_3$ is described by the map d (Remark 3). Notice that the six cliques of Γ (in red color) constitute a six-entries string of the form (4.2). Then the two colored hexagons are obtained by a polygonal enlargement (Proposition 4.2) and the pink (resp., grey) one corresponds to the clique w (resp., v). Their green colored cliques are precisely those obtained by replacing w (resp., v) by (4.4) (resp., (4.2)). The blue cliques are those belonging to $\mathfrak{C}(\partial \widetilde{P}_3) \setminus \mathfrak{C}(\partial P_3)$ but not to the enlarged perimeter $\widetilde{\Gamma}$. The whole $\widetilde{P}_3$ is portrayed in Figure 4; the first hyperbolic case ($n = 4$) is displayed in Figure 6.

5. FREE BURNSIDE GROUPS

We review here the definition and some finiteness results about free Burnside groups.

³The result, obtained by computer algebra software, is too large to be displayed here.

FIGURE 3. Cliques enumeration algorithm for $n = 3$.

Definition 5.1. The variety Bur_n of equation $x^n = 1$ is called the *variety of Burnside groups* (of order n) [14, 8].

Regard Bur_n as a category.

Lemma 5.1. For any set S of rank m there is a free object over S in Bur_n .

Proof. Let $\mathbb{F}_m$ be the free group on S and $\mathbb{F}_m^n$ the (normal) subgroup generated by its n^{th} powers. Then

$$(5.1) \quad B(m, n) := \frac{\mathbb{F}_m}{\mathbb{F}_m^n}$$

is the desired free object (over S , understood as a subset of $B(m, n)$). Indeed, if $B \in \text{Bur}_n$ and $f : S \rightarrow B$ is a map then there is a unique group homomorphism $\tilde{f} : B(m, n) \rightarrow B$ extending f . $\square$

Definition 5.2. The free object of Bur_n , with m generators, is called the *free Burnside group* of order n with m generators and denoted by $B(m, n)$.

Corollary 5.0.3. S is a Borel-free set of generators of $B(m, n)$.

Proof. The abelian group $A := \bigoplus_{s \in S} \langle s \rangle$ is an object of $\mathbf{Bur}_n$: hence, the map $f : S \rightarrow A$ which acts on S as the identity can be extended to a group homomorphism $\tilde{f} : B(m, n) \rightarrow A$.

Now, let $s \in S$ and observe that the subgroup $\langle s \rangle \leq B(m, n)$ cannot be identical. Indeed, if $\langle s \rangle = 1$ then it is impossible to extend the map $g_s : S \rightarrow \mathbb{Z}_n$ with $g_s(s) := 1$ and $g_s(s') := 0$, $s' \in S \setminus \{s\}$, to a group homomorphism, thus contradicting freedom of $B(m, n)$ over S (notice that $\mathbb{Z}_n$ is a 1-generator object of $\mathbf{Bur}_n$).

It follows that $\tilde{f}|_{\langle s \rangle}$ is injective, i.e., $\ker \tilde{f}|_{\langle s \rangle} = 1$. Let $s' \neq s$ and observe that $\tilde{f}|_{\langle s \rangle}$ maps the subgroup $\langle s \rangle \cap \langle s' \rangle \leq B(m, n)$ in the subgroup $\langle s \rangle \cap \langle s' \rangle$ of A , which is, by the definition of A , identical. In other words, $\langle s \rangle \cap \langle s' \rangle$ is a subgroup of the identical subgroup $\ker \tilde{f}|_{\langle s \rangle}$ and as such it must be identical as well. $\square$

5.1. Finiteness of $B(2, 3)$: an algebraic proof. The departing point to check finiteness of $B(2, n)$ is the expression

$$(5.2) \quad w = x^{a_1} y^{b_1} x^{a_2} y^{b_2} \dots x^{a_l} y^{b_l}$$

of an its generic element w . In one form or another, all existing algorithms to attack the Burnside problem entail manipulating the right-hand side of (5.2) in order to move all the occurrences of x (resp., y), say, on the first (resp., second) leftmost position. Performing this by brute-force inevitably brings about a surfeit of commutators:

$$(5.3) \quad w = x^{\sum a_i} y^{\sum b_i} \cdot (\text{binary commutators} \cdot \text{ternary commutators} \cdot \dots)$$

Seemingly (5.3) makes (5.2) no better at all: nevertheless, it reveals the role of nil-potency of $B(2, n)$. Indeed, if $B(2, n)$ is nilpotent then its class sets an upper bound to the multiplicity of commutators appearing in (5.3). For instance, the fact that $B(m, 2)$ is abelian (proved by Burnside in 1902 [5]), i.e., nilpotent of class 1, implies that no commutators whatsoever appear in (5.3). In 1933 Levi and Van der Waerden independently proved that $B(m, 3)$ is nilpotent of class 3, hence expression (5.3) contains commutators of multiplicity not greater than 3.

When $m = 2$ the nil-potency class of $B := B(2, 3)$ is 2, i.e., B' is central in B (and therefore B is meta-abelian). This means that all the commutators arising from the above manipulation of (5.2) can be collected in its rightmost position. In order to better appreciate the geometrical finiteness result later on (Section 7) we review here the classical proof.

Let G be a group of exponent three; in particular, $g^{-1} = g^2$ for any $g \in G$.

Proposition 5.1 ([15]). *Let $g, h \in G$. Then*

$$(5.4) \quad [g, h^2] = [h, g].$$

Proof. Observe that $1 = (gh)^3 = ghghgh$ implies

$$(5.5) \quad ghg = h^{-1}g^{-1}h^{-1} = h^2g^2h^2, \quad g, h \in G.$$

Equation (5.6) below is an identity since its both sides coincide with the same element $h^2gh^2g^2h^2$:

$$(5.6) \quad h^2(gh^2g)gh^2 = h^2g^2(g^2h^2g^2)h^2.$$

Hence, by applying (5.5) to the parenthesized expressions in (5.6) one obtains

$$h^2(hg^2h)gh^2 = h^2g^2(hgh)h^2,$$

i.e., $g^2hgh^2 = h^2g^2hg$, which is precisely $[g, h^2] = [h, g]$, and (5.4) holds true. $\square$

Corollary 5.1.1. *Every subgroup of G generated by two elements has a central derived subgroup.*

Proof. Let $H := \langle h, g \rangle$ be a two-generators subgroup. Relation (5.4) from Proposition 5.1 immediately gives

$$(5.7) \quad [g, h^2] = [g, h]^2,$$

$$(5.8) \quad [g^2, h] = [g, h]^2,$$

$$(5.9) \quad [g^2, h^2] = [g, h],$$

showing that H' is generated by $[g, h]$. Then, combining (5.7) with the general group-theoretic identity $[g, h^2] = [h, g]^{h^2}$ one obtains $[h, g] = [h, g]^{h^2}$, i.e.,

$$[h, g, h^2] = 1.$$

Similarly, from (5.8) one proves that

$$[h, g, g^2] = 1.$$

Hence, the subgroup generated by h^2 and g^2 , which is H again, commutes with H' . $\square$

Lemma 5.2 ([8, 12]). *Any element $w \in B$ can be uniquely written as*

$$(5.10) \quad w = x^a y^b [x, y]^c, \quad a, b, c \in \{0, 1, 2\}.$$

In particular, $|B| = 27$.

Proof. B' is central by Corollary 5.1.1. In particular, the normal subgroup B' is abelian and the sequence

$$\mathbb{Z}_3 \equiv B' = \langle [x, y] \rangle \longrightarrow B \longrightarrow \frac{B}{B'} = \langle xB', yB' \rangle \cong \mathbb{Z}_3^2$$

is exact, showing (5.10). $\square$

In modern classification of groups can be found precisely 5 groups of order p^3 and only two of them are non-abelian: they have exponent p or p^2 , with p odd prime number [20]. Hence, for $p = 3$, $B(2, 3)$ is the non-abelian one with exponent 3, which is also the unique non-abelian metabelian group with two generators and exponent 3 possessing a cyclic derived subgroup, and, as such, presented as

$$B(2, 3) = \langle x, y \mid x^3 = y^3 = [x, y, x] = [x, y, y] = 1 \rangle.$$

5.2. Finiteness of $B(2, 4)$: an algebraic proof. First results about finiteness of $B(2, 4)$ were given already by Burnside in his 1902 paper [5] in which he claims that $B(2, 4)$ is finite of order $\leq 2^{12}$. Next, in 1940 Sanov [22] proved that $B(m, 4)$ is finite using the following Lemma 5.3.

Lemma 5.3. [17] *Let B be a group of exponent 4, $D \leq B$ an its finite subgroup and $c \in B$ an element such that $c^2 \in D$ and $\langle c, D \rangle = B$. Then B is finite.*

Proof. Put $d := |D|$. Observe that since $B = \langle c, D \rangle$ every element $b \in B$ can be written in the form

$$(5.11) \quad b = P_1 c Q_1 c P_2 c Q_2 c \dots P_s c Q_s c P_{s+1} c Q_{s+1},$$

where $P_i, Q_i \in D$ for $1 \leq i \leq s+1$ and none of them is identical except possibly P_1 or Q_{s+1} . It is enough to show that in the expression (5.11) of b the number $2s+1$ of occurrences of the factor c is bounded by a fixed number. More precisely, we will show that $s \leq d$ by applying to the above representation of b a transformation which reduces the occurrences of the factor c if $s > d$.

Notice that $c^3 = c^{-1}$ (since $c \in B$) and that for any $R \in D$ one has $(R^{-1}c^{-1})^4 = 1$, so that

$$(5.12) \quad cRc = R^{-1}c^{-1}R^{-1}c^{-1}R^{-1} = R^{-1}c \cdot c^2R^{-1}c \cdot c^2R^{-1} = R^{-1}c\tilde{R}c\tilde{R},$$

where $\tilde{R} = c^2R^{-1} \in D$. Now, let use (5.12) to transform the representation (5.11) as follows,

$$\begin{aligned} & \dots cP_{l+k-1}cQ_{l+k-1}cP_{l+k}(cQ_{l+k}c) \dots \\ & \dots cP_{l+k-1}cQ_{l+k-1}(cP_{l+k}Q_{l+k}^{-1}c)UcU \dots \\ & \dots cP_{l+k-1}(cQ_{l+k-1}Q_{l+k}P_{l+k}^{-1}c)VcWcU \dots \\ & \dots c(P_{l+k-1}P_{l+k}Q_{l+k}^{-1}Q_{l+k-1}^{-1})cXcYcWcU \dots, \end{aligned}$$

where $U, \dots, Y \in D$. In this way we do not change the number of factors c , but just obtain a new representation of b in which two subsequent elements are separated by an element $S_{l,k}$ (or $S_{l,k}^{-1}$) of D defined as

$$S_{l,k} = P_l P_{l+1} \dots P_{l+k} Q_{l+k}^{-1} \dots Q_l^{-1},$$

where $l > 0, k \geq 0, l + k \leq s$. On the other hand, notice that any $S_{l,k} \in D$ could be extended to $S_{1,k}$, $k = 0, \dots, s-1$. Therefore if s is greater than the order d of D then at least two elements, say $S_{1,k}$ and $S_{1,l}$ must be equal, i.e.,

$$P_1 \dots P_{k+1} Q_{k+1}^{-1} \dots Q_1^{-1} = P_1 \dots P_{l+1} Q_{l+1}^{-1} \dots Q_1^{-1}.$$

Assuming $k < l$ we obtain

$$(5.13) \quad S_{k+2, l-k-1} = P_{k+2} \dots P_{l+1} Q_{l+1}^{-1} \dots Q_{k+2}^{-1} = 1.$$

Now to finish the proof it is just enough to use (5.13) in the appropriate representation of b ,

$$\dots Q_{k+1} c S_{k+2, l-k-1} c P_{l+1} \dots = \dots Q_{k+1} \underbrace{c^2}_{\in D} P_{l+1} \dots$$

□

Then, the finiteness of $B(2, 4)$ is straightforward. The group $\langle x, y^2 \rangle$, its subgroup $\langle x \rangle$ and the element y^2 fulfill the hypotheses of Lemma 5.3: hence, $\langle x, y^2 \rangle$ is finite. Then apply again Lemma 5.3 to the group $B(2, 4)$, its subgroup $\langle x, y^2 \rangle$ and the element y , and obtain the desired result.

6. THE KERNEL SUBGROUP K_n

Let $K_n := \langle D(n, n, n)^n \rangle$ be the subgroup generated by n^{th} powers of elements of $D(n, n, n)$.

Lemma 6.1 ([7, 24]). *The short sequence*

$$(6.1) \quad 0 \longrightarrow K_n \longrightarrow D(n, n, n) \longrightarrow B(2, n) \longrightarrow 0$$

is exact.

Proof. Just compare the presentation (4.1) of $D(n, n, n)$ with the definition (5.1) of $B(2, n)$ (see Lemma 5.1). □

Results from Section 4 give a geometrical flavor to sequence (6.1).

Lemma 6.2. *K_2 is trivial, K_3 is the infinite abelian subgroup generated by the cubes of the three translations xy^2 , y^2x and xyx , while K_n is an (infinite) not-abelian group for all $n > 3$.*

Proof. $D(2, 2, 2)$ is abelian of exponent 2 so $K_2 = 1$. As a subgroup of rigid motions of $\mathbb{R}^2$ (Theorem 4.1), any element $d \in D(3, 3, 3)$ can be written as rt , where r is a rotation and t a translation. It follows that $d^3 = r^3 t r^2 t r t$, i.e., d^3 is a composition of translations. Hence, $K_3 \leq \mathbb{R}^2$ is the lattice made of the centers of a regular hexagonal tiling of thrice the size as the basic hexagon of $\mathcal{P}_3$ and as such it is generated by the three vectors xy^2 , y^2x and xyx (see Fig. 4).

When $n > 3$, Corollary 4.2.2 shows that $d(3)^n$ and $d(2n-3)^n$ are aperiodic non commuting elements of K_n , which is then infinite and not abelian. □

One of the main difficulties on describing K_n for $n > 3$ is the lack of decomposition into rotations and translations (which make little sense in hyperbolic geometry). The clique enumeration algorithm will help to enumerate K_n in the hyperbolic case as well.

Corollary 6.0.2 (Kernel enumeration algorithm). *There exists a recursively defined surjection $k : \mathbb{N}_0 \rightarrow K_n$.*

Proof. For any $j \in \mathbb{N}$ define

$$\tilde{N}_j := \bigcup_{a_1 + \dots + a_l = j, l \leq j} N_{a_1} \times N_{a_2} \times \dots \times N_{a_l}$$

where the N_j 's are the same as the clique enumeration algorithm (Corollary 4.2.1). Observe that $\tilde{N}_1 = N_1$ while $\tilde{N}_2$ is the union of N_2 and N_1^2 , and so on.

Being finite each $\tilde{N}_j$ can be understood as a subset of $\mathbb{N}_0$. Furthermore, we assume that $\tilde{N}_1, \tilde{N}_2, \dots, \tilde{N}_j, \dots$ is a sequence of subsequent interval covering the whole $\mathbb{N}_0$.

Set $k \equiv d$ on $\tilde{N}_1$. Let now $i \in \tilde{N}_j$ and regard it as an l -tuple, i.e., $i \equiv (i_1, \dots, i_l)$, with $i_1 \in N_{a_1}, \dots, i_l \in N_{a_l}$. Set

$$k(i) := d(i_1)^n d(i_2)^n \dots d(i_l)^n.$$

This allows to define k on the whole $\mathbb{N}_0$.

Take now an element $k = d_1^n d_2^n \dots d_l^n \in K_n$. By the clique enumeration algorithm (Corollary 4.2.1) there exists a unique l -tuple $i_1, \dots, i_l$ such that $d_1 = d(i_1), \dots, d_l = d(i_l)$ and there exist uniques $a_1, \dots, a_l$ such that $i_1 \in N_{a_1}, \dots, i_l \in N_{a_l}$. Hence, $k \in k(\tilde{N}_j)$, where $j = a_1 + \dots + a_l$, i.e., the map k is surjective. □

Before analyzing the hyperbolic case it is convenient to show the finiteness of $B(2, 3)$ in this geometric framework.

6.1. Finiteness of $B(2, 3)$: a geometric proof. Sequence 6.1 for $n = 3$ (see Lemma 6.1) gives immediately finiteness of $B(2, 3)$. Figure 4 shows the output of the cliques enumeration algorithm (Corollary 4.2.1) stopped at the first enlargement of the basic hexagon.

According to Proposition 4.2 there are exactly 30 cliques of which 6 are pairwise identified (in Figure 4 the colored edges are just two copies of the same clique) giving the correct order of $B(2, 3)$. These 27 cliques can be easily put in correspondence with the elements of $B(2, 3)$ described in Lemma 5.2.

Table 1 displays a list of identities which follow from the properties (5.7), (5.8) and (5.9). Hence, the elements labeling the cliques in Figure 4 can be replaced by triplets of numbers ranging in $\{0, 1, 2\}$ corresponding to the powers a, b, c in (5.10). The result is displayed in Figure 5.

TABLE 1. List of the 18 nontrivial correspondences between the cliques of $T(B(2, 3), \{x, y\})$ obtained by the cliques enumeration algorithm (Corollary 4.2.1) and the elements of $B(2, 3)$ expressed as in Lemma 5.2. The remaining 9 trivial correspondences are precisely those with $c = 0$.


$\mathfrak{C}(T(B(2, 3), \{x, y\}))$	$B(2, 3)$	(a, b, c)
xyx	$x^2y[x, y]^2$	$(2, 1, 2)$
yx	$xy[x, y]^2$	$(1, 1, 2)$
yx^2	$x^2y[x, y]$	$(2, 1, 1)$
y^2x^2yx	$[x, y]^2$	$(0, 0, 2)$
$y^2x^2yx^2$	$x[x, y]^2$	$(1, 0, 2)$
y^2x^2y	$x^2[x, y]^2$	$(2, 0, 2)$
xyx^2	$y[x, y]$	$(0, 1, 1)$
y^2x	$xy^2[x, y]$	$(1, 2, 1)$
y^2x^2	$x^2y^2[x, y]^2$	$(2, 2, 2)$
xy^2x^2	$y^2[x, y]$	$(0, 2, 1)$
$(y^2x)^2x$	$y[x, y]^2$	$(0, 1, 2)$
xyx^2y	$y^2[x, y]^2$	$(0, 2, 2)$
$(y^2x^2y^2)^2$	$xy^2[x, y]^2$	$(1, 2, 2)$
xyx^2y^2	$[x, y]$	$(0, 0, 1)$
$(yx^2)^2xy$	$x^2[x, y]$	$(2, 0, 1)$
xy^2x	$x^2y^2[x, y]$	$(2, 2, 1)$
y^2xy^2	$xy[x, y]$	$(1, 1, 1)$
y^2xy	$x[x, y]$	$(1, 0, 1)$

Observe that the 7 hexagons in Figure 4 correspond precisely to a fundamental domain of K_3 which, by Lemma 6.2 is generated by the cubes of the three translations yx^2 , y^2x and xyx . Such a fundamental domain is the intersection of the three stripes determined by the three pairs of parallel lines (portrayed in the same colors in Figure 4). Within each pair of parallel lines one line is determined from the other by the cube of a translation above.

In the hyperbolic plane almost all sides of P_n and its enlargements are mutually parallel, so that compactness of fundamental domains is not taken for granted.


7. REPRESENTING THE GRAPH OF $B(2, n)$ ON A QUOTIENT OF A CONSTANT-CURVATURE SURFACES

As a discrete subgroup of the isometry group of $\mathbb{S}$ the group K_n gives rise to the quotient surface $\frac{\mathbb{S}}{K_n}$; as a group of symmetries of $\mathcal{P}_n$, K_n gives rise to a quotient tiling $\frac{\mathcal{P}_n}{K_n}$; as a group of transformations of the 2-colors graph $\partial\mathcal{P}_n$ it



Proposition 7.1. *The quotient group $\frac{D(n,n,n)}{K_n}$ is equivariantly identified with the set of cliques of the 2-colors graph $\partial\left(\frac{P_n}{K_n}\right)$ determined by the quotient $2n$ -gonal tiling $\frac{P_n}{K_n}$ of the quotient surface $\frac{\mathbb{S}}{K_n}$.*

Fix a point p in $\mathbb{H}$, e.g., the imaginary unit. For any $i \in \mathbb{N}_0$ we denote by Π_i the half-plane of $\mathbb{H}$ which contains i and is delimited by the geodesic passing through the middle point of the geodesic segment joining p and $k(i)(p)$, where

FIGURE 5. The colored graph of $B(2, 3)$ labeled by classical elements.

k is the recursive map describing K_n constructed in Corollary 6.0.2. Then the polygon

$$Q_i := \bigcap_{a=1}^i \Pi_i$$


contains a fundamental domain of K_n in $\mathbb{H}$.

Corollary 7.1.1. *There exists an algorithm which, if stops, returns a closed polygon Q containing a fundamental domain of K_n .*

Proof. Just observe that closedness of each Q_i can be checked algorithmically. $\square$


Authors are currently working on a computer–algebra implementation of the algorithm suggested in Corollary 7.1.1 in the case $n = 4$. It must be stressed that its convenience with respect to the existing coset enumeration algorithms is still to be established, especially for larger n .

Nevertheless, this approach based on colored graphs enlightens the “true reason” why a Burnside group with two generators might be infinite: it is because of the peculiarity of hyperbolic geometry, namely the existence of non–compact polygons of arbitrarily many sides.


 FIGURE 6. The colored graph of von Dyck group $D(4, 4, 4)$.

7.1. Higher-order von Dyck groups? Generalizing the result of Proposition 7.1 is by no means straightforward: it would require defining a higher-order analog of the exact sequence (6.1) involving the m -generators free Burnside group $B(m, n)$ (see (5.1)) and, eventually, a group playing the role of the von Dyck group (4.1) in the context of tilings of higher-dimensional surfaces. As an additional difficulty, the key result of Lemma 1.2 requires the surjectivity of π_* which, for $m > 2$, cannot be taken for granted (see Remark 2). So, a lot of work must be done in this direction.

The suspicion that the graph of $B(m, n)$ may be living on a three-dimensional surface is confirmed by the following toy model concerning the (finite, abelian) free Burnside group $B(3, 2) = \mathbb{Z}_2^3$. Since by Corollary 5.0.3 $B(3, 2)$ admits the Borel-free set of generators $\{x, y, z\}$, the 3-colors graph can be easily constructed and the result is displayed in Figure 7 where $H := \langle x \rangle$, $K := \langle y \rangle$ and $L := \langle z \rangle$

FIGURE 7. The graph of $B(3, 2)$.

and edges of the same color are identified. Interestingly enough, after identifications are carried out one gets a cuboctahedron (Fig. 8) though the original groups act on it by reflections (instead of rotations as in the cases with $m = 2$).

REFERENCES

- [1] Muhammad A. Albar and Walid M. Al-Hamdan. The triangle groups. *Rend. Sem. Mat. Univ. Padova*, 89:103–111, 1993.
- [2] Alan F. Beardon. *The geometry of discrete groups*, volume 91 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. Corrected reprint of the 1983 original.
- [3] Max Born and Emil Wolf. *Principles of optics: Electromagnetic theory of propagation, interference and diffraction of light*. With contributions by A. B. Bhatia, P. C. Clemmow, D. Gabor, A. R. Stokes, A. M. Taylor, P.




FIGURE 8. The cuboctahedron associated with of $B(3, 2)$.

- A. Wayman and W. L. Wilcock. Third revised edition. Pergamon Press, Oxford, 1965.
- [4] F. Buekenhout, editor. *Handbook of incidence geometry*. North-Holland, Amsterdam, 1995. Buildings and foundations.
 - [5] W Burnside. On an unsettled question in the theory of discontinuous groups. *Quart.J.Math.*, (33):230–238, 1902.
 - [6] Reinhard Diestel. *Graph theory*, volume 173 of *Graduate Texts in Mathematics*. Springer, Heidelberg, fourth edition, 2010.
 - [7] Matthew Farrelly. The burnside group $b(3; 2)$ as a two-relator quotient of $c_3 c_3$, 2007.
 - [8] Nelly Fazio, Kevin Iga, Antonio Nicolosi, Ludovic Perret, and William E. Skeith III. Hardness of learning problems over burnside groups of exponent 3. *Cryptology ePrint Archive*, Report 2011/398, 2011.
 - [9] Michael Giudici, Geoffrey Pearce, and Cheryl E. Praeger. Basic coset geometries. *J. Algebraic Combin.*, 36(4):561–594, 2012.
 - [10] Chris Godsil and Gordon Royle. *Algebraic graph theory*, volume 207 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2001.
 - [11] Branko Grünbaum and G. C. Shephard. *Tilings and patterns*. A Series of Books in the Mathematical Sciences. W. H. Freeman and Company, New

- York, 1989. An introduction.
- [12] Narain Gupta. On groups in which every element has finite order. *Amer. Math. Monthly*, 96(4):297–308, 1989.
 - [13] George Havas and M. F. Newman. Application of computers to questions like those of Burnside. In *Burnside groups (Proc. Workshop, Univ. Bielefeld, Bielefeld, 1977)*, volume 806 of *Lecture Notes in Math.*, pages 211–230. Springer, Berlin, 1980.
 - [14] L. G. Kovács. Varieties of groups and Burnside’s problem. *Bull. Amer. Math. Soc.*, 74:599–601, 1968.
 - [15] Friedrich Levi and B.L. Waerden. ber eine besondere klasse von gruppen. *Abhandlungen aus dem Mathematischen Seminar der Universitt Hamburg*, 9:154–158, 1933.
 - [16] R. Duncan Luce and Albert D. Perry. A method of matrix analysis of group structure. *Psychometrika*, 14:95–116, 1949.
 - [17] Wilhelm Magnus, Abraham Karrass, and Donald Solitar. *Combinatorial group theory*. Dover Publications Inc., Mineola, NY, second edition, 2004. Presentations of groups in terms of generators and relations.
 - [18] Antonio Pasini. Geometries and chamber systems. *Sūrikaiseikikenkyūsho Kōkyūroku*, (867):28–52, 1994. Finite groups and related topics (Japanese) (Kyoto, 1993).
 - [19] A. Pothén and S. Toledo. Elimination structures in scientific computing. In *Handbook on Data Structures and Applications*, pages Chapter 59, 29 pp. Dinesh Mehta and Sartaj Sahni, editors, Chapman and Hall /CRC, 2004.
 - [20] Derek J. S. Robinson. *A course in the theory of groups*, volume 80 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1996.
 - [21] Mark A. Ronan. Coverings and automorphisms of chamber systems. *European J. Combin.*, 1(3):259–269, 1980.
 - [22] I. N. Sanov. Solution of Burnside’s problem for exponent 4. *Leningrad State Univ. Annals [Uchenye Zapiski] Math. Ser.*, 10:166–170, 1940.
 - [23] Tits. Gomtries polydriques et groupes simples. *Atti 2a Riunione Groupem. Math. Express. Lat. Firenze*, pages 66–88, 1962.
 - [24] A. M. Vinogradov. Why is space three-dimensional and how may groups be seen? *Acta Appl. Math.*, 5(2):169–180, 1986.
 - [25] K. S. Wang and L. C. Grove. A note on Von Dyck groups. *Exposition. Math.*, 9(3):285–288, 1991.

* MATHEMATICAL INSTITUTE IN OPAVA
 SILESIAN UNIVERSITY IN OPAVA
 NA RYBNICKU 626/1
 746 01 OPAVA
 CZECH REPUBLIC
 E-mail address: Giovanni.Moreno@math.slu.cz

** DEPARTMENT OF MATHEMATICS
SALERNO UNIVERSITY
VIA PONTE DON MELILLO
84084 SALERNO
ITALY
E-mail address: `mstypa@unisa.it`